



## Free choosability of outerplanar graphs

Yves Aubry, Jean-Christophe Godin, Olivier Togni

### ► To cite this version:

Yves Aubry, Jean-Christophe Godin, Olivier Togni. Free choosability of outerplanar graphs. *Graphs and Combinatorics*, 2016, 32 (3), pp.851-859. 10.1007/s00373-015-1625-3 . hal-00957298

**HAL Id: hal-00957298**

**<https://hal.science/hal-00957298>**

Submitted on 10 Mar 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# FREE CHOOSABILITY OF THE CYCLE

YVES AUBRY, JEAN-CHRISTOPHE GODIN AND OLIVIER TOGNI

**ABSTRACT.** A graph  $G$  is free  $(a, b)$ -choosable if for any vertex  $v$  with  $b$  colors assigned and for any list of colors of size  $a$  associated with each vertex  $u \neq v$ , the coloring can be completed by choosing for  $u$  a subset of  $b$  colors such that adjacent vertices are colored with disjoint color sets. In this note, a necessary and sufficient condition for a cycle to be free  $(a, b)$ -choosable is given. As a corollary, some choosability results are derived for graphs in which cycles are connected by a tree structure.

## 1. INTRODUCTION

For a graph  $G$ , we denote its vertex set by  $V(G)$  and edge set by  $E(G)$ . A color-list  $L$  of a graph  $G$  is an assignment of lists of integers (colors) to the vertices of  $G$ . For an integer  $a$ , a  $a$ -color-list  $L$  of  $G$  is a color-list such that  $|L(v)| = a$  for any  $v \in V(G)$ .

In 1996, Voigt considered the following problem: let  $G$  be a graph and  $L$  a color-list and assume that an arbitrary vertex  $v \in V(G)$  is precolored by a color  $f \in L(v)$ . Under which hypothesis is it always possible to complete this precoloring to a proper color-list coloring ? This question leads to the concept of free choosability introduced by Voigt [8].

Formally, for a graph  $G$ , integers  $a, b$  and a  $a$ -color-list  $L$  of  $G$ , an  $(L, b)$ -coloring of  $G$  is a mapping  $c$  that associates to each vertex  $u$  a subset  $c(u)$  of  $L(u)$  such that  $|c(u)| = b$  and  $c(u) \cap c(v) = \emptyset$  for any edge  $uv \in E(G)$ . The graph  $G$  is  $(a, b)$ -choosable if for any  $a$ -color-list  $L$  of  $G$ , there exists an  $(L, b)$ -coloring. Moreover,  $G$  is free  $(a, b)$ -choosable if for any  $a$ -color-list  $L$ , any vertex  $v$  and any set  $c_0 \subset L(v)$  of  $b$  colors, there exists an  $(L, b)$ -coloring  $c$  such that  $c(v) = c_0$ .

As shown by Voigt [8], there are examples of graphs  $G$  that are  $(a, b)$ -choosable but not free  $(a, b)$ -choosable. Some related recent results concern defective free choosability of planar graphs [6]. We investigate, in the next section, the free-choosability of the first interesting case, namely the cycle. We derive a necessary and sufficient condition for a cycle to be free  $(a, b)$ -choosable (Theorem 4). In order to get a concise statement, we introduce the free-choice ratio of a graph, in the same way that Alon, Tuza and Voigt [1] introduced the choice ratio (which is equal to the so-called fractional chromatic number).

For any real  $x$ , let  $\text{FCH}(x)$  be the set of graphs  $G$  which are free  $(a, b)$ -choosable for all  $a, b$  such that  $\frac{a}{b} \geq x$ :

---

*Date:* March 10, 2014.

*2010 Mathematics Subject Classification.* 05C15, 05C38, 05C72.

*Key words and phrases.* Coloring, Choosability, Free choosability, Cycles.

$$\text{FCH}(x) = \left\{ G \mid \forall \frac{a}{b} \geq x, G \text{ is free } (a, b)\text{-choosable} \right\}.$$

Moreover, we can define the free-choice ratio  $\text{fchr}(G)$  of a graph  $G$  by:

$$\text{fchr}(G) := \inf \left\{ \frac{a}{b} \mid G \text{ is free } (a, b)\text{-choosable} \right\}.$$

**Remark 1.** *Erdős, Rubin and Taylor have asked [3] the following question: If  $G$  is  $(a, b)$ -choosable, and  $\frac{c}{d} > \frac{a}{b}$ , does it imply that  $G$  is  $(c, d)$ -choosable? Gutner and Tarsi have shown [5] that the answer is negative in general. If we consider the analogue question for the free choosability, then Theorem 4 implies that it is true for the cycle.*

The path  $P_{n+1}$  of length  $n$  is the graph with vertex set  $V = \{v_0, v_1, \dots, v_n\}$  and edge set  $E = \bigcup_{i=0}^{n-1} \{v_i v_{i+1}\}$ . The cycle  $C_n$  of length  $n$  is the graph with vertex set  $V = \{v_0, \dots, v_{n-1}\}$  and edge set  $E = \bigcup_{i=0}^{n-1} \{v_i v_{i+1(\text{mod } n)}\}$ . To simplify the notation, for a color-list  $L$  of  $P_n$  or  $C_n$ , we let  $L(i)$  denote  $L(v_i)$  and  $c(i)$  denote  $c(v_i)$ .

The notion of waterfall color-list was introduced in [2] to obtain choosability results on the weighted path and then used to prove the  $(5m, 2m)$ -choosability of triangle-free induced subgraphs of the triangular lattice. We recall one of the results from [2] that will be used in this note, with the function  $\text{Even}$  being defined for any real  $x$  by:  $\text{Even}(x)$  is the smallest even integer  $p$  such that  $p \geq x$ .

**Proposition 2** ([2]). *Let  $L$  be a color-list of  $P_{n+1}$  such that  $|L(0)| = |L(n)| = b$ , and  $|L(i)| = a = 2b + e$  for all  $i \in \{1, \dots, n-1\}$  (with  $e > 0$ ).*

*If  $n \geq \text{Even}\left(\frac{2b}{e}\right)$  then  $P_{n+1}$  is  $(L, b)$ -colorable.*

For example, let  $P_{n+1}$  be the path of length  $n$  with a color-list  $L$  such that  $|L(0)| = |L(n)| = 4$ , and  $|L(i)| = 9$  for all  $i \in \{1, \dots, n-1\}$ . Then the previous proposition tells us that we can find an  $(L, 4)$ -coloring of  $P_{n+1}$  whenever  $n \geq 8$ . In other words, if  $n \geq 8$ , we can choose 4 colors on each vertex such that adjacent vertices receive disjoint colors sets. If  $|L(i)| = 11$  for all  $i \in \{1, \dots, n-1\}$ , then  $P_{n+1}$  is  $(L, 4)$ -colorable whenever  $n \geq 4$ . On the other side, there are color-lists  $L$  for which  $P_{n+1}$  is not  $(L, b)$ -colorable

## 2. FREE CHOOSABILITY OF THE CYCLE

We begin with a negative result for the even-length cycle:

**Lemma 3.** *For any integers  $a, b, p$  such that  $p \geq 2$ , and  $\frac{a}{b} < 2 + \frac{1}{p}$ , the cycle  $C_{2p}$  is not free  $(a, b)$ -choosable.*

*Proof.* We construct a counterexample for the free-choosability of  $C_{2p}$ : let  $L$  be the  $a$ -color-list of  $C_{2p}$  such that

$$L(i) = \begin{cases} \{1, \dots, a\}, & \text{if } i \in \{0, 1\}; \\ \{\frac{i-1}{2}a + 1, \dots, \frac{i-1}{2}a + a\}, & \text{if } i \neq 2p-1 \text{ is odd}; \\ \{b + \frac{i-2}{2}a + 1, \dots, b + (\frac{i-2}{2} + 1)a\}, & \text{if } i \text{ is even and } i \neq 0; \\ \{1, \dots, b, 1 + (p-1)a, \dots, 1 + pa - b - 1\}, & \text{if } i = 2p-1. \end{cases}$$




FIGURE 1. The cycle  $C_6$ , along with a 9-color-list  $L$  for which there is no  $(L, 4)$ -coloring  $c$  such that  $c(v_0) = \{1, 2, 3, 4\}$ .

If we choose  $c_0 = \{1, \dots, b\} \subset L(0)$ , we can check that there does not exist an  $(L, b)$ -coloring of  $C_{2p}$  such that  $c(0) = c_0$ , so  $C_{2p}$  is not free  $(a, b)$ -choosable. See Figure 1 for an illustration when  $p = 3$ ,  $a = 9$  and  $b = 4$ .  $\square$

Now, if  $\lfloor x \rfloor$  denotes the greatest integer less than or equal to the real  $x$ , we can state:

**Theorem 4.** *For the cycle  $C_n$  of length  $n$ ,*

$$C_n \in \text{FCH} \left( 2 + \left\lfloor \frac{n}{2} \right\rfloor^{-1} \right).$$

Moreover, we have:

$$\text{fchr}(C_n) = 2 + \left\lfloor \frac{n}{2} \right\rfloor^{-1}.$$

*Proof.* Let  $a, b$  be two integers such that  $a/b \geq 2 + \lfloor \frac{n}{2} \rfloor^{-1}$ . Let  $L$  be a  $a$ -color-list of  $C_n$ . Without loss of generality, we can suppose that  $v_0$  is the vertex chosen for the free-choosability and  $c_0 \subset L(v_0)$  has  $b$  elements. It remains to construct an  $(L, b)$ -coloring  $c$  of  $C_n$  such that  $c(v_0) = c_0$ . Hence we have to construct an  $(L', b)$ -coloring  $c$  of  $P_{n+1}$  such that  $L'(0) = L'(n) = L_0$  and for all  $i \in \{1, \dots, n-1\}$ ,  $L'(i) = L(v_i)$ . We have  $|L'(0)| = |L'(n)| = b$  and for all  $i \in \{1, \dots, n-1\}$ ,  $|L'(i)| = a$ . Since  $a/b \geq 2 + \lfloor \frac{n}{2} \rfloor^{-1}$  and  $e = a - 2b$ , we get  $e/b \geq \lfloor \frac{n}{2} \rfloor^{-1}$  hence  $n \geq \text{Even}(2b/e)$ . Using Proposition 2, we get:

$$C_n \in \text{FCH} \left( 2 + \left\lfloor \frac{n}{2} \right\rfloor^{-1} \right).$$

Hence, we have that  $\text{fchr}(C_n) \leq 2 + \lfloor \frac{n}{2} \rfloor^{-1}$ . Moreover, let us prove that  $M = 2 + \lfloor \frac{n}{2} \rfloor^{-1}$  is reached. For  $n$  odd, Voigt has proved [9] that the choice ratio  $\text{chr}(C_n)$  of a cycle of odd length  $n$  is exactly  $M$ . Hence  $\text{fchr}(C_n) \geq \text{chr}(C_n) = M$ , and the result is proved. For  $n$  even, Lemma 3 asserts that  $C_n$  is not free  $(a, b)$ -choosable for  $\frac{a}{b} < 2 + \lfloor \frac{n}{2} \rfloor^{-1}$ .  $\square$

**Remark 5.** *In particular, the previous theorem implies that if  $b, e, n$  are integers such that  $n \geq \text{Even}(\frac{2b}{e})$ , then the cycle  $C_n$  of length  $n$  is free  $(2b + e, b)$ -choosable.*

In order to extend the result to other graphs than cycles, the following simple proposition will be useful:

**Proposition 6.** *Let  $a, b$  be integers with  $a \geq 2b$ . Let  $G$  be a graph and  $G_v$  be the graph obtained by adding a leave  $v$  to any vertex of  $G$ . Then  $G$  is free  $(a, b)$ -choosable if and only if  $G_v$  is free  $(a, b)$ -choosable.*

*Proof.* Since the "only if" part holds trivially, let us prove the "if" part. Assume  $G$  is free  $(a, b)$ -choosable and let  $L$  be a  $a$ -color-list of  $G$ . Let  $v$  be a new vertex and let  $G_v$  be the graph obtained from  $G$  by adding the edge  $uv$ , for some  $u \in V(G)$ . Then any  $(L, b)$ -coloring  $c$  of  $G$  can be extended to an  $(L, b)$ -coloring of  $G_v$  by giving to  $v$   $b$  colors from  $L(v) \setminus c(u)$  ( $|L(v) \setminus c(u)| \geq b$  since  $a \geq 2b$ ). If  $v$  is colored with  $b$  colors from its list, then, since  $G$  is free  $(a, b)$ -choosable, the coloring can be extended to an  $(L, b)$ -coloring of  $G_v$  by first choosing for  $u$  a set of  $b$  colors from  $L(u) \setminus c(v)$ .  $\square$

Starting from a single edge and applying inductively Proposition 6 allows to obtain the following corollary:

**Corollary 7.** *Let  $T$  be a tree of order  $n \geq 2$ . Then*

$$T \in \text{FCH}(2).$$

Now, we can state the following:

**Proposition 8.** *If  $G$  is a unicyclic graph with girth  $g$ , then*

$$G \in \text{FCH} \left( 2 + \left\lfloor \frac{g}{2} \right\rfloor^{-1} \right).$$

*Proof.* Let  $a, b$  be two integers such that  $a/b \geq 2 + \lfloor \frac{g}{2} \rfloor^{-1}$ ,  $L$  be a  $a$ -color-list of  $G$ ,  $C = v_1, \dots, v_g$  be the unique cycle (of length  $g$ ) of  $G$  and  $T_i$ ,  $i \in \{1, \dots, g\}$ , be the subtree of  $G$  rooted at vertex  $v_i$  of  $C$ .

Let  $v$  be the vertex chosen for the free choosability and let  $c_0 \subset L(v)$  be a set of cardinality  $b$ . If  $v \in C$ , then by virtue of Theorem 4, there exists an  $(L, b)$ -coloring  $c$  of  $C$  such that  $c(v) = c_0$ . This coloring can be easily extended to the whole graph by coloring the vertices of each tree  $T_i$  thanks to Corollary 7. If  $v \in T_i$  for some  $i$ ,  $1 \leq i \leq g$ , then Corollary 7 asserts that the coloring can be extended to  $T_i$ . Then color  $C$  starting at vertex  $v_i$  by using Theorem 4. Finally, complete it by coloring each tree  $T_j$ ,  $1 \leq j \neq i \leq g$ .  $\square$

### 3. APPLICATIONS

As an example to the possible use of the results from Section 2, we begin with determining the free choosability of a binocular graph, i.e. two cycles linked by a path.

For integers  $m, n$  and  $p$  such that  $m, n \geq 3$  and  $p \geq 0$ , the *binocular graph*  $BG(m, n, p)$  is the disjoint union of an  $m$ -cycle  $u_0, u_1, \dots, u_{m-1}$  and of an  $n$ -cycle  $v_0, \dots, v_{n-1}$  with vertices  $u_0$  and  $v_0$  linked by a path of length  $p$  given by  $u_0, x_1, \dots, x_{p-1}, v_0$ . Note that if  $p = 0$ , then  $u_0$  and  $v_0$  are the same vertex.

**Proposition 9.** *For any  $m \geq 3$ ,  $n \geq 3$  and  $p \geq 0$ ,*

$$BG(m, n, p) \in \text{FCH} \left( 2 + \left\lfloor \frac{\min(m, n)}{2} \right\rfloor^{-1} \right).$$




FIGURE 2. A tree of cycles (on the left), and the associated tree obtained by collapsing cycles (on the right).

*Proof.* Assume without loss of generality that  $m \geq n$  and let  $a, b$  be integers such that  $a/b \geq 2 + \lfloor \frac{n}{2} \rfloor^{-1}$ . Let  $L$  be a  $a$ -color-list of  $BG(m, n, p)$ . Let  $y$  be the vertex chosen for the free choosability and let  $c_0 \subset L(y)$  be a set of cardinality  $b$ . If  $y$  lies on the  $m$ -cycle, then by virtue of Theorem 4, there exists an  $(L, b)$ -coloring  $c$  of the  $m$ -cycle such that  $c(y) = c_0$ . By Corollary 7, this coloring can be extended to the vertices of the path. Now, it remains to color the vertices of the  $n$ -cycle, with  $v_0$  being already colored. This can be done thanks to Theorem 4. If  $y$  lies on the  $n$ -cycle, the argument is similar. If  $y \in \{x_1, \dots, x_{p-1}\}$ , then the coloring can be extended to the whole path and the coloring of the  $m$ -cycle and  $n$ -cycle can be completed thanks to Theorem 4.  $\square$

This method can be extended to prove similar results on graphs with more than two cycles, connected by a tree structure.

Define a *tree of cycles* to be a graph  $G$  such that all its cycles are disjoint and collapsing all vertices of each cycle of  $G$  produces a tree.

**Corollary 10.** *Any tree of cycles of girth  $g$  is in  $FCH(2 + \lfloor \frac{g}{2} \rfloor^{-1})$ .*

#### 4. ALGORITHMIC CONSIDERATIONS

Let  $n \geq 3$  be an integer and let  $a, b$  be two integers such that  $a/b \geq 2 + \lfloor \frac{n}{2} \rfloor^{-1}$ . Let  $L$  be a  $a$ -color-list of  $C_n$ .

As defined in [2], a *waterfall* list  $L$  of a path  $P_{n+1}$  of length  $n$  is a list  $L$  such that for all  $i, j \in \{0, \dots, n\}$  with  $|i - j| \geq 2$ , we have  $L(i) \cap L(j) = \emptyset$ . Let  $m = |\cup_{i=0}^n L(i)|$  be the total number of colors of the color-list  $L$ .

The algorithm behind the proof of Proposition 2 consists in three steps: first, the transformation of the list  $L$  into a waterfall list  $L'$  by renaming some colors; second, the construction of the  $(L', b)$ -coloring by coloring vertices from 0 to  $n - 1$ , giving to vertex  $i$  the first  $b$ -colors that are not used by the previous vertex; third, the backward transformation to obtain an  $(L, b)$ -coloring from the  $(L', b)$ -coloring by coming back to original colors and resolving color conflicts if any. It can be seen that the time complexity of the first step is  $O(mn)$ ; that of the second one is  $O(a^2n)$  and that of the third one is  $O(\max(a, b^3)n)$ . Therefore, the total running time for computing a free  $(L, b)$ -coloring of the cycle  $C_n$  is  $O(\max(m, a^2, b^3)n)$ .

## REFERENCES

- [1] N. Alon, Zs. Tuza, M. Voigt, *Choosability and fractional chromatic number*, Discrete Math. 165/166, (1997), 31-38.
- [2] Y. Aubry, J.-C. Godin and O. Togni, *Every triangle-free induced subgraph of the triangular lattice is  $(5m, 2m)$ -choosable*, Discrete Applied Math. 166, (2014), 51-58.
- [3] P. Erdős, A.L Rubin and H. Taylor, *Choosability in graphs*, Proc. West-Coast Conf. on Combinatorics, Graph Theory and Computing, Congressus Numerantium XXVI, (1979), 125-157.
- [4] J.-C. Godin, *Coloration et choisissabilité des graphes et applications*, PhD thesis (in french), Université du Sud Toulon-Var, France (2009).
- [5] S. Gutner and M. Tarsi, *Some results on  $(a:b)$ -choosability*, Discrete Math. 309, (2009), 2260-2270.
- [6] N. Li, X. Meng and H. Liu, *Free Choosability of Outerplanar Graphs*, Green Communications and Networks, Lecture Notes in Electrical Engineering, 113, (2012), 253-256.
- [7] Zs. Tuza and M. Voigt, *Every 2-choosable graph is  $(2m, m)$ -choosable*, J. Graph Theory 22, (1996), 245-252.
- [8] M. Voigt, *Choosability of planar graphs*, Discrete Math., 150, (1996), 457-460.
- [9] M. Voigt, *On list Colourings and Choosability of Graphs*, Abilitationsschrift, TU Ilmenau (1998).

INSTITUT DE MATHÉMATIQUES DE TOULON, UNIVERSITÉ DE TOULON, FRANCE, IN-  
STITUT DE MATHÉMATIQUES DE MARSEILLE, FRANCE

*E-mail address:* `yves.aubry@univ-tln.fr`, `godinjeanchri@yahoo.fr`

LABORATOIRE LE2I UMR CNRS 6306, UNIVERSITÉ DE BOURGOGNE, FRANCE

*E-mail address:* `olivier.togni@u-bourgogne.fr`